

AD632236

The Statistics Center
RUTGERS-THE STATE UNIVERSITY

TWO-STAGE CHAIN SAMPLING INSPECTION PLANS
WITH DIFFERENT SAMPLE SIZES IN THE TWO STAGES

by

K. S. Stephens

and

H. F. Dodge

GPO PRICE \$ _____

CFSTI PRICE(S) \$ _____

Hard copy (HC) 2.00

Microfiche (MF) 1.50

653 July 65

TECHNICAL REPORT NO. N-23

April, 1966

Research supported by the Army, Navy, Air Force, and NASA under Office of Naval Research Contract No. Nonr 404(18) Task No. NR 042-241 with Rutgers-The State University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

N66 32133

FACILITY FORM 802

(ACCESSION NUMBER)
45
(PAGES)
CR-76626
(NASA CR OR TMX OR AD NUMBER)
AD-632236

(THRU)
1
(CODE)
19
(CATEGORY)

rwfNwo

TWO-STAGE CHAIN SAMPLING INSPECTION PLANS
WITH DIFFERENT SAMPLE SIZES IN THE TWO STAGES

by

K. S. Stephens*

and

H. F. Dodge

INTRODUCTION

In previous reports^{1,2,3,4} a generalized family of two-stage chain sampling inspection plans was presented. Conditions for their use were also discussed. This report contains a further generalization of these plans with particular attention to the specification of the sample sizes used in the different stages of the criteria. Previous plans specify the use of the same sample size in both stages. The plans presented here call for the use of different sample sizes in the two stages.

While this modification results in a variable amount of inspection, it has certain advantages. In fact, it seems logical to require a larger sample during initial start-up and following rejected lots, for, in practice, it often happens that defective lots occur in bunches. Under these conditions if a defective lot is found, as suggested by a rejection, the succeeding lots warrant closer inspection. In principle, the inspection procedures for both the military standard plans (MIL-STD-105D⁵), and the continuous sampling plans (such as CSP-1⁶) also incorporate intensified inspections during periods of possibly excessive defectiveness. MIL-STD-105D provides for "Tightened Inspection" when lot rejections are too closely spaced, while CSP-1 requires

* This report is based in part on work being done in preparation of a doctoral thesis at Rutgers-The State University.

100% inspection (to the extent of clearing i consecutive units) following the occurrence of a defective during sampling.

Comparisons of the new plans with those having the same parameters but a single sample size indicate that improved discrimination is achieved by this two-sample-size procedure. The evaluation of the new plans is carried out using the framework of the theory described in a previous report.³

GENERAL PLAN AND OPERATING PROCEDURE

The general plan and operating procedure are the same as described in the above-mentioned previous reports. The parameters designating the number of samples over which cumulation takes place in the different stages i.e., k_1 and k_2 , and the parameters designating the allowable number of defectives in the associated cumulative results i.e., C_1 and C_2 , remain the same. The parameter designating the sample size i.e., n , is now defined separately for each of the two stages--namely n_1 and n_2 .

For purposes of illustration and completeness, a modified schematic of the operating procedure given in the previous reports is shown in Fig. 1.

The following designations are used in the schematic to describe the operation of the sampling procedure:

d_i --the number of defectives in the i th sample.

D_i --the cumulative number of defectives at the i th sample, with cumulation performed as shown.

Here the definition of D_i differs from that used in the earlier report.¹




Fig. 1 Flow Chart of Operations, Two-Stage Chain Sampling Plan
with Different n , i.e. n_1 and n_2

DESIGNATION OF PLANS

It is of course possible to assign a large number of different values to the six parameters which make up a plan. To facilitate discussion the general system of designation defined before will again be used, with $\text{ChSP}(n_1, n_2) - C_1, C_2$ designating the "Chain Sampling Plan with Different n , i.e. n_1 and n_2 " with cumulative-result acceptance numbers C_1 and C_2 :

<u>Parameters</u>						<u>Designation</u>
n_1 ,	n_2 ;	k_1 ,	k_2 ;	C_1 ,	C_2	
n_1 ,	n_2 ;	k_1 ,	k_2 ;	0,	1	$\text{ChSP}(n_1, n_2) - 0, 1$
n_1 ,	n_2 ;	k_1 ,	k_2 ;	0,	2	$\text{ChSP}(n_1, n_2) - 0, 2$
n_1 ,	n_2 ;	k_1 ,	k_2 ;	1,	2	$\text{ChSP}(n_1, n_2) - 1, 2$
n_1 ,	n_2 ;	k_1 ,	k_2 ;	0,	3	$\text{ChSP}(n_1, n_2) - 0, 3$
n_1 ,	n_2 ;	k_1 ,	k_2 ;	1,	3	$\text{ChSP}(n_1, n_2) - 1, 3$
etc.						etc.

In referring to a specific plan, the values of the parameters in the above order will be enclosed in parentheses e.g., (20, 10; 1, 2; 0, 2) designates the $\text{ChSP}(n_1, n_2) - 0, 2$ plan having parameters: $n_1 = 20$, $n_2 = 10$; $k_1 = 1$, $k_2 = 2$; $C_1 = 0$, $C_2 = 2$. On occasion it will be convenient to abridge the notation to the four basic parameters, when n_1 and n_2 are clearly implied.

EVALUATING OPERATING CHARACTERISTICS

The operating characteristics of the present procedure are readily evaluated using the Markov chains developed for evaluating the ChSP procedures of the previous reports. The only necessary modification is to make each transition probability a function of the parameters n_1 and n_2 , depending on whether the

transition involves the first or second stage of the criteria. In general, the transition probabilities from those states not involving an "R" (denoting a rejection) will always be functions of n_2 only, since these states are reached only while in the second stage of the criteria. The transition probabilities from state "R" will always be functions of n_1 only, since a rejection requires a return to the first stage of the criteria. The transition probabilities from other states involving an R, e.g., (R0, R1, R2, R00, R01, R10, R02, R02, R000, etc.) will be functions of n_1 or n_2 depending on the parameter k_1 . When the number of sample outcomes contained in a state involving a rejection (R) are equal to or greater than $k_1 + 1$, transitions from such states will be functions of n_2 . But when the number of sample outcomes contained in a state involving a rejection (R) are less than $k_1 + 1$, transitions from these states will be functions of n_1 . To illustrate the situation for states involving an R other than the state "R" itself, the following examples are given.

Example 1

Let $k_1 = 1$, $k_2 = 4$; $C_1 = 0$, $C_2 = 2$.

States involving an R other than the state "R" itself are:

$\left. \begin{array}{l} \text{R00} \\ \text{R01} \\ \text{R02} \end{array} \right\}$ number of sample outcomes contained in each state equal 3,
 R0 - number of sample outcomes contained in this state equal 2.

The number of sample outcomes contained in each of these states $\geq k_1 + 1 = 2$.

Hence transitions from these states are functions of n_2 , since passage to the second stage of the criteria takes place with the occurrence of zero defectives following a rejection (i.e., $k_1 = 1$). The possible transitions from the above states in accordance with the given acceptance criteria are as follows:

$P_{\text{R00,000}}$ = Probability of zero defectives in n_2

$P_{R00,001}$ = Probability of one defective in n_2

$P_{R00,002}$ = Probability of two defectives in n_2

$P_{R00,R}$ = Probability of three or more defectives in n_2

$P_{R01,010}$ = Probability of zero defectives in n_2

$P_{R01,011}$ = Probability of zero defectives in n_2

$P_{R01,R}$ = Probability of two or more defectives in n_2

$P_{R02,020}$ = Probability of zero defectives in n_2

$P_{R02,R}$ = Probability of one or more defectives in n_2

$P_{R0,R00}$ = Probability of zero defectives in n_2

$P_{R0,R01}$ = Probability of one defective in n_2

$P_{R0,R02}$ = Probability of two defectives in n_2

$P_{R0,R}$ = Probability of three or more defectives in n_2

Now note that for $k_1 = 2$, $k_2 = 4$; $C_1 = 0$, $C_2 = 2$, only two of the above states are admissible--namely R00 and R0, i.e., R01 and R02 exceed the $C_1 = 0$ criterion for $k_1 = 2$ samples following a rejection.

For the first of the admissible states involving a rejection i.e., R00, the number of sample outcomes contained in the state = $k_1 + 1 = 3$.

Hence transitions from this state are functions of n_2 and are the same as in the above example.

However, the number of sample outcomes contained in the state $R0 < k_1 + 1 = 3$.

Hence transitions from this state are functions of n_1 since two (i.e., k_1) acceptable sample outcomes must follow a rejection before passing to the second stage of the criteria.

The possible transitions from this state in accordance with the given acceptance criteria are as follows:

$P_{RO,ROO}$ = Probability of zero defectives in n_1 .

$P_{RO,R}$ = Probability of one or more defectives in n_1 .

Example 2

Let $k_1 = 3$, $k_2 = 4$; $C_1 = 0$, $C_2 = 2$.

For this case, states ROO and RO are again admissible. However, the number of sample outcomes contained in both states ROO and RO $< k_1 + 1 = 4$. Hence transitions from both states are functions of n_1 .

The possible transitions are as follows:

$P_{ROO,000}$ = Probability of zero defectives in n_1

$P_{ROO,R}$ = Probability of one or more defectives in n_1

$P_{RO,ROO}$ = Probability of zero defectives in n_1

$P_{RO,R}$ = Probability of one or more defectives in n_1

Previously (in the earlier reports) the transition probabilities, p_{ij} , were labeled: P_d , $d = 0, 1, \dots, C_2$ and defined accordingly,

P_d = Probability of d defectives in a sample of n

For this report the transition probabilities, p_{ij} are labeled:

P_{d,n_s} , $d = 0, 1, \dots, C_2$; $s = 1, 2$ and are defined as,

P_{d,n_s} = Probability of d defectives in a sample of n_s .

Making use of this notation, the transition probability matrices for a number of Markov chains are given below. These are Markov chains for a selected group of the sets of parameters which have been evaluated. The following nine plans are illustrated:

Plan	$k_1,$	$k_2;$	$C_1,$	C_2
(1)	1,	2;	0,	1
(2)	1,	3;	0,	1
(3)	2,	3;	0,	1
(4)	1,	2;	0,	2
(5)	1,	3;	0,	2
(6)	2,	3;	1,	2
(7)	1,	5;	0,	2
(8)	1,	2;	0,	3
(9)	2,	3;	0,	3

Algebraic solutions for P_a , the proportion of lots expected to be accepted, are given for plans (1), (2), (3), (4), and (8) by solving the Markov chains for the limiting probability of the rejection, "R", state.

Plan (1): 1,2; 0,1

		State at ith trial		
		0	1	R
State at (i-1)st trial	0	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
	1	P_{0,n_2}	.	$1 - P_{0,n_2}$
	R	P_{0,n_1}	.	$1 - P_{0,n_1}$

Fig. 2. Markov Chain for ChSP(n_1, n_2)-0,1 Plan: ($k_1, k_2; C_1, C_2 = 1, 2; 0, 1$)

Proceeding as in previous reports^{3,4}, the solution of the limiting probability of state "R" from which P_a is obtained is as follows:

$$P_0 = P_{0,n_2} P_0 + P_{0,n_2} P_1 + P_{0,n_1} P_R \quad (1)$$

$$P_1 = P_{1,n_2} P_0 \quad (2)$$

$$P_0 + P_1 + P_R = 1 \quad (3)$$

Using (2) in (1) produces the following equations:

$$P_0 = \frac{P_{0,n_1} P_R}{1 - P_{0,n_2} - P_{0,n_2} P_{1,n_2}} \quad (4)$$

$$P_1 = \frac{P_{0,n_1} P_{1,n_2} P_R}{1 - P_{0,n_2} - P_{0,n_2} P_{1,n_2}} \quad (5)$$

Using (4) and (5) in (3) gives the result for P_R

$$P_R = \frac{1 - P_{0,n_2} - P_{0,n_2} P_{1,n_2}}{1 + P_{0,n_1} - P_{0,n_2} + P_{0,n_1} P_{1,n_2} - P_{0,n_2} P_{1,n_2}} \quad (6)$$

$$P_a = 1 - P_R = \frac{P_{0,n_1} + P_{0,n_1} P_{1,n_2}}{1 + P_{0,n_1} - P_{0,n_2} + P_{0,n_1} P_{1,n_2} - P_{0,n_2} P_{1,n_2}} \quad (7)$$

It is of interest to check the agreement of this result with the special case of $n_1 = n_2 = n$, for which the parameters being considered are those of a ChSP-0,1 plan¹ or more specifically, the ChSP-1 plan⁷ with $i = 1$.

With $P_{0,n_1} = P_{0,n_2} = P_0$, and $P_{1,n_2} = P_1$, (7) reduces to,

$$P_a = P_0 + P_0 P_1, \text{ which is the expected result.}$$

Plan (2): 1,3; 0,1

		State at ith trial				
		00	01	10	R0	R
State at (i-1)st trial	00	$P_{0,n_2} P_{1,n_2}$.	.	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
	01	.	.	P_{0,n_2}	.	$1 - P_{0,n_2}$
	10	P_{0,n_2}	.	.	.	$1 - P_{0,n_2}$
	R0	$P_{0,n_2} P_{1,n_2}$.	.	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
	R	.	.	.	P_{0,n_1}	$1 - P_{0,n_1}$

Fig. 3. Markov Chain for ChSP(n_1, n_2)-0,1 Plan: (k_1, k_2 ; $C_1, C_2 = 1, 3; 0, 1$)

$$P_{00} = P_{0,n_2} P_{00} + P_{0,n_2} P_{10} + P_{0,n_2} P_{R0} \quad (8)$$

$$P_{01} = P_{1,n_2} P_{00} + P_{1,n_2} P_{R0} \quad (9)$$

$$P_{10} = P_{0,n_2} P_{01} \quad (10)$$

$$P_{R0} = P_{0,n_1} P_R \quad (11)$$

$$P_{00} + P_{01} + P_{10} + P_{R0} + P_R = 1 \quad (12)$$

Combining equations as before,

$$P_R = \frac{1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2}}{1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1} + P_{0,n_1} P_{1,n_2} + P_{0,n_1} P_{0,n_2} P_{1,n_2}} \quad (13)$$

$$P_a = \frac{P_{0,n_1} + P_{0,n_1} P_{1,n_2} + P_{0,n_1} P_{0,n_2} P_{1,n_2}}{1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1} + P_{0,n_1} P_{1,n_2} + P_{0,n_1} P_{0,n_2} P_{1,n_2}} \quad (14)$$

The result can also be checked against the special case for $n_1 = n_2 = n$.

With $P_{0,n_1} = P_{0,n_2} = P_0$, and $P_{1,n_2} = P_1$, (14) reduces to,

$$P_a = \frac{P_0 + P_0 P_1 + P_0^2 P_1}{1 + P_0 P_1}, \text{ which is the result}$$

obtained from (1), page 8 of Reference 1 when $k_1 = 1$, and

$k_2 = 3$.

Plan (3): 2,3; 0,1

		State at ith trial				
		00	01	10	R0	R
State at (i-1)st trial	00	P_{0,n_2}	P_{1,n_2}	.	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
	01	.	.	P_{0,n_2}	.	$1 - P_{0,n_2}$
	10	P_{0,n_2}	.	.	.	$1 - P_{0,n_2}$
	R0	P_{0,n_1}	.	.	.	$1 - P_{0,n_1}$
	R	.	.	.	P_{0,n_1}	$1 - P_{0,n_1}$

Fig. 4. Markov Chain for ChSP(n_1, n_2)-0,1 Plan: (k_1, k_2 ; $C_1, C_2 = 2, 3; 0, 1$)

$$P_{00} = P_{0,n_2} P_{00} + P_{0,n_2} P_{10} + P_{0,n_1} P_{R0} \quad (15)$$

$$P_{01} = P_{1,n_2} P_{00} \quad (16)$$

$$P_{10} = P_{0,n_2} P_{01} \quad (17)$$

$$P_{R0} = P_{0,n_1} P_R \quad (18)$$

$$P_{00} + P_{01} + P_{10} + P_{R0} + P_R = 1 \quad (19)$$

Combining equations as before,

$$P_R = (1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2}) / (1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1} - P_{0,n_1} P_{0,n_2} - P_{0,n_1} P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1}^2 + P_{0,n_1}^2 P_{1,n_2} + P_{0,n_1}^2 P_{0,n_2} P_{1,n_2}) \quad (20)$$

$$\begin{aligned}
 P_a = 1 - P_R = & (P_{0,n_1} - P_{0,n_1} P_{0,n_2} - P_{0,n_1} P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1}^2 P_{0,n_2} + P_{0,n_1}^2 P_{1,n_2} \\
 & + P_{0,n_1}^2 P_{0,n_2} P_{1,n_2}) / (1 - P_{0,n_2} - P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1} \\
 & - P_{0,n_1} P_{0,n_2} - P_{0,n_1} P_{0,n_2}^2 P_{1,n_2} + P_{0,n_1}^2 P_{0,n_2} + P_{0,n_1}^2 P_{1,n_2} + P_{0,n_1}^2 P_{1,n_2} \\
 & + P_{0,n_1}^2 P_{0,n_2} P_{1,n_2}) \quad (21)
 \end{aligned}$$

Again letting $P_{0,n_1} = P_{0,n_2} = P_0$, and $P_{1,n_2} = P_1$, (21) reduces to,

$P_a = P_0 + P_0^2 P_1$, which is the result for the associated ChSP-0,1 plan with $n_1 = n_2 = n$.

Plan (4): 1,2; 0,2

		State at ith trial			
		0	1	2	R
State at (i-1)st trial	0	P_{0,n_2}	P_{1,n_2}	P_{2,n_2}	$1 - \sum_{i=0}^2 P_{i,n_2}$
	1	P_{0,n_2}	P_{1,n_2}	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
	2	P_{0,n_2}	.	.	$1 - P_{0,n_2}$
	R	P_{0,n_1}	.	.	$1 - P_{0,n_1}$

Fig. 5. Markov Chain for ChSP(n_1, n_2)-0,2 Plan: (k_1, k_2 ; $C_1, C_2 = 1, 2; 0, 2$)

$$P_0 = P_{0,n_2} P_0 + P_{0,n_2} P_1 + P_{0,n_2} P_2 + P_{0,n_1} P_R \quad (22)$$

$$P_1 = P_{1,n_2} P_0 + P_{1,n_2} P_1 \quad (23)$$

$$P_2 = P_{2,n_2} P_0 \quad (24)$$

$$P_0 + P_1 + P_2 + P_R = 1 \quad (25)$$

These combine to give,

$$\begin{aligned} P_a = & (P_{0,n_1} + P_{0,n_1} P_{2,n_2} - P_{0,n_1} P_{1,n_2} P_{2,n_2}) / (1 + P_{0,n_1} \\ & + P_{0,n_1} P_{2,n_2} - P_{0,n_1} P_{1,n_2} P_{2,n_2} - P_{0,n_2} - P_{1,n_2} \\ & - P_{0,n_2} P_{2,n_2} + P_{0,n_2} P_{1,n_2} P_{2,n_2}) \end{aligned} \quad (26)$$

With $P_{0,n_1} = P_{0,n_2} = P_0$, $P_{1,n_2} = P_1$ and $P_{2,n_2} = P_2$, (26) reduces to,

$$P_a = \frac{P_0 + P_0 P_2 - P_0 P_1 P_2}{1 - P_1}, \text{ which agrees}$$

with (19), page 13, Reference 3.

Plan (5): 1,3; 0,2

		State at ith trial							
		00	01	10	11	02	20	R0	R
State at (i-1)st trial	00	P_{0,n_2}	P_{1,n_2}	.	.	P_{2,n_2}	.	.	$1 - \sum_{i=0}^2 P_{i,n_2}$
	01	.	.	P_{0,n_2}	P_{1,n_2}	.	.	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
	10	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
	11	.	.	P_{0,n_2}	$1 - P_{0,n_2}$
	02	P_{0,n_2}	.	$1 - P_{0,n_2}$
	20	P_{0,n_2}	$1 - P_{0,n_2}$
	R0	P_{0,n_2}	P_{1,n_2}	.	.	P_{2,n_2}	.	.	$1 - \sum_{i=0}^2 P_{i,n_2}$
	R	P_{0,n_1}	$1 - P_{0,n_1}$

Fig. 6. Markov Chain for ChSP(n_1, n_2)-0,2 Plan: (k_1, k_2 ; $C_1, C_2 = 1, 3; 0, 2$)

Plan (6): 2,3; 1,2

State at ith trial

	00	01	10	11	02	20	R0	R1	R
00	P_{0,n_2}	P_{1,n_2}	.	.	P_{2,n_2}	.	.	.	$1 - \sum_{i=0}^2 P_{i,n_2}$
01	.	.	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
10	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
11	.	.	P_{0,n_2}	$1 - P_{0,n_2}$
02	P_{0,n_2}	.	.	$1 - P_{0,n_2}$
20	P_{0,n_2}	$1 - P_{0,n_2}$
R0	P_{0,n_1}	P_{1,n_1}	$1 - \sum_{i=0}^1 P_{i,n_1}$
R1	.	.	P_{0,n_1}	$1 - P_{0,n_1}$
R	P_{0,n_1}	P_{1,n_1}	$1 - \sum_{i=0}^1 P_{i,n_1}$

Fig. 7. Markov Chain for ChSP(n_1, n_2)-1,2 Plan: (k_1, k_2 ; $C_1, C_2 = 2, 3; 1, 2$)

Plan (7): 1,5; 0,2

State at i th trial

	0000	0001	0010	0100	1000	1100	1010	1001	0110	0101	0011	0002	0020	0200	2000	R000	R001	R010	R011	R020	R00	R01	R02	R0	R
0000	P_{0,n_2}	P_{1,n_2}	P_{2,n_2}	$1 - \frac{2}{150} P_{1,n_2}$	R
0001	.	P_{0,n_2}	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
0010	.	.	P_{0,n_2}	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
0100	.	.	.	P_{0,n_2}	.	.	.	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
1000	P_{0,n_2}	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
1100	.	.	.	P_{0,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
1010	.	.	P_{0,n_2}	$1 - P_{0,n_2}$	R
1001	.	P_{0,n_2}	$1 - P_{0,n_2}$	R
0110	P_{0,n_2}	$1 - P_{0,n_2}$	R
0101	P_{0,n_2}	$1 - P_{0,n_2}$	R
0011	P_{0,n_2}	$1 - P_{0,n_2}$	R
0002	P_{0,n_2}	$1 - P_{0,n_2}$	R
0020	P_{0,n_2}	$1 - P_{0,n_2}$	R
0200	P_{0,n_2}	$1 - P_{0,n_2}$	R
2000	P_{0,n_2}	P_{0,n_2}	$1 - P_{0,n_2}$	R
R000	P_{0,n_2}	P_{1,n_2}	P_{2,n_2}	$1 - \frac{2}{150} P_{1,n_2}$	R
R001	.	P_{0,n_2}	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
R010	.	.	P_{0,n_2}	P_{1,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
R011	.	.	.	P_{0,n_2}	.	.	.	P_{0,n_2}	$1 - \frac{1}{150} P_{1,n_2}$	R
R002	P_{0,n_2}	$1 - P_{0,n_2}$	R
R020	P_{0,n_2}	$1 - P_{0,n_2}$	R
R00	P_{0,n_2}	.	$P_{0,n_2} P_{1,n_2}$	$1 - \frac{2}{150} P_{1,n_2}$	R
R01	$P_{0,n_2} P_{1,n_2}$	$1 - \frac{1}{150} P_{1,n_2}$	R
R02	$P_{0,n_2} P_{1,n_2}$	$1 - \frac{1}{150} P_{1,n_2}$	R
R0	P_{0,n_2}	$1 - P_{0,n_2}$	R
R	$P_{0,n_2} P_{1,n_2} P_{2,n_2}$.	.	$1 - \frac{2}{150} P_{1,n_2}$	R

Fig. 8. Markov Chain for ChSP(n_1, n_2)-0,2 Plan: (k_1, k_2 ; $C_1, C_2 = 1,5; 0,2$)

Plan (8): 1,2; 0,3

State at (i-1)st trial	State at ith trial				
	0	1	2	3	R
0	P_{0,n_2}	P_{1,n_2}	P_{2,n_2}	P_{3,n_2}	$1 - \sum_{i=0}^3 P_{i,n_2}$
1	P_{0,n_2}	P_{1,n_2}	P_{2,n_2}	.	$1 - \sum_{i=0}^2 P_{i,n_2}$
2	P_{0,n_2}	P_{1,n_2}	.	.	$1 - \sum_{i=0}^1 P_{i,n_2}$
3	P_{0,n_2}	.	.	.	$1 - P_{0,n_2}$
R	P_{0,n_1}	.	.	.	$1 - P_{0,n_1}$

Fig. 9. Markov Chain for ChSP(n_1, n_2)-0,3 Plan: (k_1, k_2 ; $C_1, C_2 = 1, 2; 0, 3$)

$$P_0 = P_{0,n_2} P_0 + P_{0,n_2} P_1 + P_{0,n_2} P_2 + P_{0,n_2} P_3 + P_{0,n_1} P_R \quad (27)$$

$$P_1 = P_{1,n_2} P_0 + P_{1,n_2} P_1 + P_{1,n_2} P_2 \quad (28)$$

$$P_2 = P_{2,n_2} P_0 + P_{2,n_2} P_1 \quad (29)$$

$$P_3 = P_{3,n_2} P_0 \quad (30)$$

$$P_0 + P_1 + P_2 + P_3 + P_R = 1 \quad (31)$$

These combine to give,

$$P_a = [P_{0,n_1} (1 + P_{2,n_2} + P_{3,n_2} - P_{1,n_2} P_{3,n_2} - P_{1,n_2} P_{2,n_2} P_{3,n_2})] / [1 + P_{0,n_1} (1 + P_{2,n_2} + P_{3,n_2} - P_{1,n_2} P_{3,n_2} - P_{1,n_2} P_{2,n_2} P_{3,n_2}) - P_{0,n_2} (1 + P_{2,n_2} + P_{3,n_2} - P_{1,n_2} P_{3,n_2} - P_{1,n_2} P_{2,n_2} P_{3,n_2}) - P_{0,n_2} P_{1,n_2} P_{2,n_2} P_{3,n_2} - P_{0,n_2} P_{1,n_2} P_{2,n_2}] \quad (32)$$

With $P_{0,n_1} = P_{0,n_2} = P_0$, $P_{1,n_2} = P_1$, $P_{2,n_2} = P_2$, and $P_{3,n_2} = P_3$, (32)

reduces to,

$$P_a = \frac{P_0 + P_0 P_2 + P_0 P_3 - P_0 P_1 P_3 - P_0 P_1 P_2 P_3}{1 - P_1 - P_1 P_2} \quad \text{which}$$

agrees with (9), page 4, Reference 4.

Plan (9): 2,3; 0,3

		State at ith trial											
		00	01	10	11	02	20	21	12	03	30	RO	R
State at (i-1)st trial	00	P_{0,n_2}	P_{1,n_2}	.	.	P_{2,n_2}	.	.	.	P_{3,n_2}	.	.	$1 - \sum_{i=0}^3 P_{i,n_2}$
	01	.	.	P_{0,n_2}	P_{1,n_2}	.	.	.	P_{2,n_2}	.	.	.	$1 - \sum_{i=0}^2 P_{i,n_2}$
	10	P_{0,n_2}	P_{1,n_2}	.	.	P_{2,n_2}	$1 - \sum_{i=0}^2 P_{i,n_2}$
	11	.	.	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
	02	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
	20	P_{0,n_2}	P_{1,n_2}	$1 - \sum_{i=0}^1 P_{i,n_2}$
	21	.	.	P_{0,n_2}	$1 - P_{0,n_2}$
	12	P_{0,n_2}	$1 - P_{0,n_2}$
	03	P_{0,n_2}	.	$1 - P_{0,n_2}$
	30	P_{0,n_2}	$1 - P_{0,n_2}$
	RO	P_{0,n_1}	$1 - P_{0,n_1}$
	R	P_{0,n_1}	$1 - P_{0,n_1}$

Fig. 10. Markov Chain for ChSP(n_1, n_2)-0,3 Plan: (k_1, k_2 ; $C_1, C_2 = 2,3; 0,3$)

By the use of the theory of Markov chains, the operating characteristics for a given value of fraction defective p over the effective range of p have been carried out for a large number of ChSP(n_1, n_2) plans. Plans with $C_1, C_2 = 0, 1; 0, 2; 0, 3$ and $1, 3; k_2 = 2, 3$ and 5 and $n_1 = 2n_2$, for $n_2 = 5, 10$, and 100 have been evaluated. OC curves for these plans are shown in Appendix A, Figs. 11 through 16.

Determination of ASN

As noted, the sampling plans being considered here result in a variable amount of sampling inspection. A measure of this amount as a function of the fraction defective is that of the average sample size or Average Sample Number (ASN) as it is called. This statistic can readily be determined for the n_1, n_2 chain sampling plans in terms of the limiting probabilities of the appropriate states.

In general the ASN is as follows:

$$ASN = n_1 P_{n_1} + n_2 P_{n_2}, \text{ where}$$

P_{n_1} = Proportion of trials (e.g. lots) for which n_1 is required.

P_{n_2} = Proportion of trials (e.g. lots) for which n_2 is required
 $= 1 - P_{n_1}$.

To illustrate with an example, consider the ChSP(n_1, n_2)-0,2 plan with parameters ($n_1, n_2; 1, 2; 0, 2$).

The possible states of the sampling process are: $0, 1, 2$ and R , representing the cumulative defectives over $k_2 - 1$ trials. n_1 is required after each rejection for one ($k_1 = 1$) trial. This occurs with probability P_R .

Hence, $P_{n_1} = P_R; P_{n_2} = 1 - P_R$, and

$$ASN = n_1 P_R + n_2(1-P_R) = n_2 + n_1 P_R - n_2 P_R.$$

To explore this more generally a somewhat more complicated example must be considered, say $ChSP(n_1, n_2) - 0, 2$ with parameters $(n_1, n_2; 2, 3, ; 0, 2)$.

The possible states are: 00, 01, 10, 11, 02, 20, R0, and R.

In this case we cannot say that n_1 will be required twice ($k_1=2$) after each rejection since rejections can occur consecutively.

However only two states can be reached following a rejection, i.e., R and R0, the 1st (i.e. R) when a second rejection results and the 2nd (i.e. R0) when an acceptance ($D \leq C_1$) results. Thus n_1 is required after each rejection and after each acceptance just preceded by a rejection.

$$\text{Hence } P_{n_1} = P_R + P_{R0}$$

$$P_{n_2} = 1 - P_R - P_{R0} = \sum_i P_i, \quad i = 00, 01, 10, 11, 02, 20$$

$$ASN = n_1(P_R + P_{R0}) + n_2(1 - P_R - P_{R0}), \text{ which can be reduced as follows,}$$

$$= n_1(P_R + P_0 P_R) + n_2(1 - P_R - P_0 P_R)$$

$$= n_2 + n_1 P_R(1 + P_0) - n_2 P_R(1 + P_0)$$

where P_0 = probability of zero defectives in a sample of n_1

with fraction defective, p .

Note that for each of the two cases considered above, when $n_1 = n_2 = n$,

$$ASN = n.$$

In general then,

for $k_1 = 0$, n_1 is not used.

$k_1 = 1$, n_1 is required after each rejection, for all k_2 ,

i.e. $k_2 \geq k_1 + 1$

$$P_{n_1} = P_R$$

$k_1 \geq 2$, n_1 is required after each rejection and after from one

to $k_1 - 1$ acceptances just preceded by a rejection.

$P_{n_1} = \sum_{j \in S} P_j$, i.e. the sum of the limiting probabilities of those states satisfying the above conditions. S corresponds to the set of all states of the associated Markov chain which involve an R and contain k_1 or less sample outcomes (counting R).
(Note that this holds for $k_1 = 1$ also, since the only state satisfying this condition is "R" itself)

$$\text{thus, ASN} = n_1 \sum_{j \in S} P_j + n_2 (1 - \sum_{j \in S} P_j) \quad (33)$$

A previous report³ contains the procedure for solving the Markov chains for the limiting probabilities of the states with particular emphasis on the "R" state so that $P_a = 1 - P_R$ is obtained. By the same procedure the limiting probabilities of the other states can be found in order to determine ASN. The states for which limiting probabilities are needed, as noted above, are those states containing an R. These are readily reduced to functions of P_R from the Markov chains by the following relation and by the nature of the construction of the states.

$$P_j = \sum_{i=1}^s p_{ij} P_i, \quad j \in S \quad (34)$$

where s = the total number of states

p_{ij} = the transition probabilities, generally

denoted by P_i , i.e. the probability of getting

i defectives in a sample of n with fraction defective, p .

e.g. for $(n_1, n_2; 2, 3; 0, 2)$

$P_{RO} = P_0 P_R$, where P_0 = the probability of getting zero defectives in a sample of n_1 with fraction defective p .

and for $(n_1, n_2; 4, 5; 1, 2)$

$P_{R0001} = P_1 P_{ROO} = P_1 P_0 P_{RO} = P_1 P_0^2 P_R$, with P_0 and P_1 similarly defined.

For the special case of $n_1 = 2n_2$ which applies to all of the specific evaluations carried out in the present report, the formulation of ASN reduces further. In this case expression (33) becomes,

$$ASN = n_2 + n_2 \sum_{j \in S} P_j \quad (35)$$

Further study has shown that for the ChSP-0,1; 0,2; 1,2; 0,3 and 1,3 plans with $k_2 = 2, 3$ and 5 which are considered in this report, five distinct formulas for ASN result. These are as follows:

For all plans with $k_1 = 1$, from $(1, 2; 0, 1)$ to $(1, 5; 1, 3)$,

$$ASN = n_2 + n_2 P_R, \quad (36)$$

For plans with $C_1, C_2 = 0, 1; 0, 2$ and $0, 3$ and $k_1 = 2$

$$ASN = n_2 + n_2(1 + P_0) P_R, \quad (37)$$

For plans with $C_1, C_2 = 0, 1; 0, 2$ and $0, 3$ and $k_1 = 4$,

$$ASN = n_2 + n_2(1 + P_0 + P_0^2 + P_0^3) P_R, \quad (38)$$

For plans with $C_1, C_2 = 1, 2$ and $1, 3$ and $k_1 = 2$,

$$ASN = n_2 + n_2(1 + P_0 + P_1) P_R \quad (39)$$

For plans with $C_1, C_2 = 1, 2$ and $1, 3$ and $k_1 = 4$,

$$ASN = n_2 + n_2 (1 + P_0 + P_0^2 + P_0^3 + 2P_0P_1 + 3P_0^2P_1) P_R, \quad (40)$$

where in all cases P_0 and P_1 denote the probabilities of 0 and 1 defectives in a sample of n_1 with fraction defective, p .

Formulas (36) through (40) have been used to obtain the results which are presented in Appendix B. A set of ASN curves for $n_1=200$, $n_2=100$ for k_1 , $k_2=1, 2$; $1, 3$; and $2, 3$ is shown in Fig. 17. Another set for $k_1, k_2=1, 5$; $2, 5$; and $4, 5$ is shown in Fig. 18. These sets correspond directly to the ChSP plans for which the OC curves are presented in Figs. 15 and 16, Appendix A. The ASN curves for the sample size combination, $n_1=200$, $n_2=100$, are shown as being representative of the effect of the ChSP parameters on this property.

DISCUSSION OF OPERATING CHARACTERISTICS

The OC curves shown in Figs. 11 through 16 of Appendix A have been arranged differently than those of the preceding reports. Three sets of two pages each are used for the different sample size combinations--Figs. 11 and 12 for $n_1=10$, $n_2=5$; Figs. 13 and 14 for $n_1=20$, $n_2=10$; and Figs. 15 and 16 for $n_1=200$, $n_2=100$. Each chart within a page contains OC curves for different k_1, k_2 combinations-- $k_1, k_2=1, 2$; $1, 3$; and $2, 3$ on the first page of a set and $k_1, k_2=1, 5$; $2, 5$; and $4, 5$ on the second page of a set. Thus the effect of changing k_1 is shown between each of the individual charts as is

the effect of changing k_2 also. The effect of different C_1 is shown within each individual chart. By this arrangement the combinations of C_1 , i.e. $C_1, C_2 = 0, 1; 0, 2$ and $1, 2; 0, 3$ and $1, 3$, of three preceding reports^{1, 3, 4} are all covered in this single report for the more general $\text{ChSP}(n_1, n_2) - C_1, C_2$ plans. The OC curves are of Type B*, based on probabilities of sampling from a process.

By comparing the OC curves of this report with those of the previous reports it can be seen that all of the OC curves presented here are tightened considerably over their constant n counterparts (where $n = n_2$ of these plans); the scales have been halved. However, within this tightened range of p values it is seen that the effects of k_1, k_2, C_1 , and C_2 are the same as for the constant n case. We need only to summarize these effects, then.

Effect of k_1 and k_2

Increasing k_1 from 1 up to $k_2 - 1$ has the effect of tightening (i.e. lowering) the OC curves, more drastically for $\text{ChSP}(n_1, n_2) - 0, C_2$ than for $\text{ChSP}(n_1, n_2) - 1, C_2$. (The $k_1 = 0$ plans are not shown due to the poor discrimination of these plans and because they are not part of the $\text{ChSP}(n_1, n_2)$ set of plans (i.e. they involve only one stage). Increasing k_2 also has the effect of tightening the OC curves. However, in the case of a fixed k_1 , say $k_1 = 1$, this tightening is accompanied by a somewhat poorer discrimination. For the plans in which $k_1 = k_2 - 1$, the effect of increasing k_2 is to tighten the OC curves considerably.

Effect of C_1 and C_2

In the $\text{ChSP} - C_1, C_2$ plans of previous reports it was noted that the effect of using a 0, 1; 0, 2; or 0, 3 chain sampling plan is to add a swelling on the

* See Reference 8, pp. 56-60.

underlying OC curve of the $c=0$, given n , single sampling plan. It was also noted that the swelling increases (particularly for low values of fraction defective, p) by going successively from a 0,1 to a 0,2 and 0,3 plan. It can be seen from the OC curves presented here that a similar effect is true for the $\text{ChSP}(n_1, n_2)-0, C_2$ plans, although here the swelling is added to the $c=0, n_1$, single sampling plan. The $\text{ChSP}(n_1, n_2)-1, C_2$ plans, like their constant n counterparts, have less discriminating OC curves.

The cost that is incurred in terms of additional inspection effort for the greatly improved discrimination and tightening of the OC curves of the $\text{ChSP}(n_1, n_2)-C_1, C_2$ plans (with $n_1=2n_2$) is provided, to a large extent, by the average sample number, ASN. As noted, sample sets of ASN curves for $n_1=200, n_2=100$ are shown in Figs. 17 and 18 of Appendix B. It can be seen from these curves that ASN is effected most significantly by the parameter k_1 (which, of course, specifies the number of samples of n_1 required after each rejection). In general, the curves tend to support the use of plans with the smaller k_1 values, i.e. $k_1=1$ or 2.

To facilitate comparison of the $\text{ChSP}(n_1, n_2)-C_1, C_2$ plans with their $\text{ChSP}-C_1, C_2$ counterparts, two additional sets of OC curves are presented in Figs. 19 and 20 of Appendix C. Fig. 19 contains OC curves for parameter sets; $k_1, k_2=1, 2; 1, 3; 2, 3$, and $1, 5; C_1, C_2=0, 1$ and $0, 2$ for $n_1, n_2=20, 10$ and for $n=10$. Fig. 20 contains OC curves for parameter sets: $k_1, k_2=1, 2, 1, 3, 2, 3$, and $2, 5; C_1, C_2=0, 2$ and $0, 3$ for $n_1, n_2=200, 100$ and for $n=100$. The curves clearly illustrate the effect of the larger sample size in the first stage, i.e. $n_1=2n_2$.

REFERENCES

1. Dodge, H. F. and K. S. Stephens, "A General Family of Chain Sampling Inspection Plans," Rutgers University Statistics Center, Technical Report No. N-20, December 30, 1964.
2. Dodge, H. F. and K. S. Stephens, "Some New Chain Sampling Inspection Plans," Annual Technical Conference Transactions, American Society for Quality Control, May 1965, pp. 8-17.
3. Stephens, K. S. and H. F. Dodge, "Chain Sampling Inspection Plans--ChSP-0,2 and ChSP-1,2," Rutgers University Statistics Center, Technical Report No. N-21, May 1965.
4. Stephens, K. S. and H. F. Dodge, "Chain Sampling Inspection Plans--ChSP-0,3 and ChSP-1,3," Rutgers University Statistics Center, Technical Report No. N-22, December 1965.
5. MIL-STD-105D, Sampling Procedures and Tables for Inspection by Attributes, U. S. Government Printing Office, Washington, D. C., 29 April 1963.
6. Dodge, H. F., "A Sampling Inspection Plan for Continuous Production, Annals of Mathematical Statistics, Vol. 14, No. 3, Sept. 1943, pp. 264-279; also Transactions of the A.S.M.E., Vol. 66, No. 2, Feb. 1944, pp. 127-133. See also: Dodge, H. F., "Sampling Plans for Continuous Production," Industrial Quality Control, Vol. IV, No. 3, Nov. 1947, pp. 5-9.
7. Dodge, H. F., "Chain Sampling Inspection Plan," Industrial Quality Control, Vol. XI, No. 4, January 1955, pp. 10-13.
8. Dodge, H. F. and H. G. Romig, Sampling Inspection Tables, John Wiley and Sons, Inc., New York, 2nd Edition, 1959.

Appendix A: Operating Characteristic Curves for ChSP(n_1, n_2)- C_1, C_2 Plans

OC curves for the following plans are presented here. Figs. 11, 12; 13, 14; and 15, 16 give OC curves for sample size combinations, $n_1, n_2=10, 5$; 20, 10; and 200, 100 respectively.

Figs. 11.1, 13.1 and 15.1 Figs. 11.2, 13.2 and 15.2 Figs. 11.3, 13.3 and 15.3
 $n_1, n_2=10, 5$; 20, 10; 200, 100 $n_1, n_2=10, 5$; 20, 10; 200, 100 $n_1, n_2=10, 5$; 20, 10; 200, 100

k_1	k_2	C_1	C_2
1	2	1	3
1	2	1	2
1	2	0	3
1	2	0	2
1	2	0	1

k_1	k_2	C_1	C_2
1	3	1	3
1	3	1	2
1	3	0	3
1	3	0	2
1	3	0	1

k_1	k_2	C_1	C_2
2	3	1	3
2	3	1	2
2	3	0	3
2	3	0	2
2	3	0	1

Figs. 12.1, 14.1 and 16.1 Figs. 12.2, 14.2 and 16.2 Figs. 12.3, 14.3 and 16.3
 $n_1, n_2=10, 5$; 20, 10; 200, 100 $n_1, n_2=10, 5$; 20, 10; 200, 100 $n_1, n_2=10, 5$; 20, 10; 200, 100

k_1	k_2	C_1	C_2
1	5	1	3
1	5	1	2
1	5	0	3
1	5	0	2
1	5	0	1

k_1	k_2	C_1	C_2
2	5	1	3
2	5	1	2
2	5	0	3
2	5	0	2
2	5	0	1

k_1	k_2	C_1	C_2
4	5	1	3
4	5	1	2
4	5	0	3
4	5	0	2
4	5	0	1




Fig. 11. OC Curves for ChSP(10,5)- C_1, C_2 Plans, $k_1, k_2 = 1,2; 1,3; 2,3$.




Fig. 12. OC Curves for ChSP(10,5)- C_1, C_2 Plans, $k_1, k_2 = 1, 5; 2, 5; 4, 5$.




Fig. 13. OC Curves for ChSP(20,10)- C_1, C_2 Plans, $k_1, k_2 = 1, 2; 1, 3; 2, 3$.






Fig. 14. OC Curves for ChSP(20,10)- C_1, C_2 Plans, $k_1, k_2 = 1, 5; 2, 5; 4, 5$.




Fig. 15. OC Curves for ChSP(200,100)- C_1, C_2 Plans, $k_1, k_2 = 1, 2; 1, 3; 2, 3$.




Fig. 16. OC Curves for ChSP(200,100)- C_1, C_2 Plans, $k_1, k_2 = 1, 5; 2, 5; 4, 5$.

Appendix B: Average Sample Number Curves for ChSP(n_1, n_2)- C_1, C_2 Plans

ASN curves for the following plans are presented here, for the sample size combination $n_1, n_2=200, 100$.

Fig. 17.1

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
1	2	1	3
1	2	1	2
1	2	0	3
1	2	0	2
1	2	0	1

Fig. 17.2

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
1	3	1	3
1	3	1	2
1	3	0	3
1	3	0	2
1	3	0	1

Fig. 17.3

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
2	3	1	3
2	3	1	2
2	3	0	3
2	3	0	2
2	3	0	1

Fig. 18.1

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
1	5	1	3
1	5	1	2
1	5	0	3
1	5	0	2
1	5	0	1

Fig. 18.2

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
2	5	1	3
2	5	1	2
2	5	0	3
2	5	0	2
2	5	0	1

Fig. 18.3

<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
4	5	1	3
4	5	1	2
4	5	0	3
4	5	0	2
4	5	0	1




Fig. 17. ASN Curves for ChSP(200,100)- C_1, C_2 Plans, $k_1, k_2 = 1, 2; 1, 3; 2, 3$.




Fig. 18. ASN Curves for ChSP(200,100)-C₁,C₂ Plans, k₁,k₂ = 1,5; 2,5; 4,5.

Appendix C: Operating Characteristic Curves for ChSP(n_1, n_2)- C_1, C_2 and ChSP- C_1, C_2 Plans.

OC curves for the following plans are presented here.

Fig. 19.1

Fig. 19.2

Fig. 19.3

Fig. 19.4

<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
10	10	1	2	0	2	10	10	1	3	0	2	10	10	2	3	0	2	10	10	1	5	0	2
		1	2	0	1			1	3	0	1			2	3	0	1			1	5	0	1
20	10	1	2	0	2	20	10	1	3	0	2	20	10	2	3	0	2	20	10	1	5	0	2
		1	2	0	1			1	3	0	1			2	3	0	1			1	5	0	1

Fig. 20.1

Fig. 20.2

Fig. 20.3

Fig. 20.4

<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>	<u>n_1</u>	<u>n_2</u>	<u>k_1</u>	<u>k_2</u>	<u>C_1</u>	<u>C_2</u>
100	100	1	2	0	3	100	100	1	3	0	3	100	100	2	3	0	3	100	100	2	5	0	3
		1	2	0	2			1	3	0	2			2	3	0	2			2	5	0	2
200	100	1	2	0	3	200	100	1	3	0	3	200	100	2	3	0	3	200	100	2	5	0	3
		1	2	0	2			1	3	0	2			2	3	0	2			2	5	0	2




Fig. 19. Comparison of OC Curves for ChSP(20,10)- C_1, C_2 Plans and ChSP- C_1, C_2 Plans with $n=10$.







Fig. 20 Comparison of OC Curves for ChSP(200,100)- C_1, C_2 Plans and ChSP- C_1, C_2 Plans with $n=100$.

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
Statistics Center; Rutgers-The State University		Unclassified	
		2b. GROUP	
3. REPORT TITLE			
Two-stage Chain Sampling Inspection Plans With Different Sample Sizes in the Two Stages			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report			
5. AUTHOR(S) (Last name, first name, initial)			
Stephens, Kenneth S. and Harold F. Dodge			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
April, 1966		38	8
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr 404(18)		Technical Report N-23	
b. PROJECT NO.			
RR 003-05-01			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
Task No. NR 042-241			
d.			
10. AVAILABILITY/LIMITATION NOTICES			
Distribution of this document is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Logistics & Mathematical Statistics Branch Office of Naval Research Washington, D.C. 20360	
13. ABSTRACT			
<p>This report presents a further generalization of the family of "two-stage" chain sampling inspection plans--the use of different sample sizes in the two stages. Plans presented in previous reports specify the use of the same sample size in both stages. Evaluation of the operating characteristics is accomplished by the Markov chain approach developed in the preceding reports. Markov chains for a number of plans are included and several algebraic solutions are developed. Since these plans involve a variable amount of sampling, an evaluation of the average sample number (ASN) is developed. OC curves are presented and discussed for plans with acceptance numbers, $C_1, C_2 = 0, 1; 0, 2, 1, 2; \text{ and } 0, 3, 1, 3$ and sample sizes, $n_1, n_2 = 10, 5; 20, 10; \text{ and } 200, 100$. ASN curves for the same plans for $n_1, n_2 = 200, 100$ are also presented. Then to compare the two-sample-size plans with plans having only one sample size, OC curves for a number of each of these types of plans are presented. Comparisons indicate that improved discrimination is achieved by the two-sample-size plans.</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Chain Sampling Sampling Inspection Plans Acceptance Sampling Cumulative-results Plans Markov Chains Operating Characteristics Average Sample Number						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.